

The role of environmental stress in fruit pigmentation

Richard V. Espley¹  | Laura Jaakola^{2,3} 

¹Department of New Cultivar Innovation, The New Zealand Institute for Plant and Food Research Ltd, Auckland, New Zealand

²Department of Arctic and Marine Biology, UiT The Arctic University of Norway, Tromsø, Norway

³Norwegian Institute of Bioeconomy Research (NIBIO), Ås, Norway

Correspondence

Richard V. Espley, Department of New Cultivar Innovation, The New Zealand Institute for Plant and Food Research Ltd, Auckland 1025, New Zealand.
Email: richard.espley@plantandfood.co.nz

Funding information

NordPlant; The New Zealand Government Ministry of Business, Innovation and Employment; The New Zealand Government Ministry of Business, Innovation and Employment's Endeavour Fund, Grant/Award Number: C11X1704

Abstract

For many fruit crops, the colour of the fruit outwardly defines its eating quality. Fruit pigments provide reproductive advantage for the plant as well as providing protection against unfavourable environmental conditions and pathogens. For consumers these colours are considered attractive and provide many of the dietary benefits derived from fruits. In the majority of species, the main pigments are either carotenoids and/or anthocyanins. They are produced in the fruit as part of the ripening process, orchestrated by phytohormones and an ensuing transcriptional cascade, culminating in pigment biosynthesis. Whilst this is a controlled developmental process, the production of pigments is also attuned to environmental conditions such as light quantity and quality, availability of water and ambient temperature. If these factors intensify to stress levels, fruit tissues respond by increasing (or ceasing) pigment production. In many cases, if the stress is not severe, this can have a positive outcome for fruit quality. Here, we focus on the principal environmental factors (light, temperature and water) that can influence fruit colour.

KEYWORDS

anthocyanin, carotenoid, climate, environmental stress, fruit, pigment, transcription

1 | BACKGROUND

The colour of fruit is generally associated with reproduction, specifically a means to promote seed dispersal by making the fruit visually attractive to herbivores/omnivores. Colour, therefore, is a key component of fruit development and intrinsically related to the ripening process. Fruit colours are derived from 'secondary (or specialised) metabolite' pathways. In evolutionary terms, these pathways in flowering plants and the breadth of their products have grown in number and complexity, in response to environmental and pathogenic stimuli and to adaptive changes for pollinator attraction. These colour pigments are largely derived from either the carotenoid and/or flavonoid pathways.

Colour helps to define the attractiveness and quality of fruits and strongly influences consumer demand. Hence, for many years, improvements or changes to fruit colour have been a major target

of breeding programmes across a range of crops (Figure 1). Additionally, growers routinely implement a number of on-orchard management techniques to enhance fruit colour development (Wang et al., 2020). In this review, we will consider the impact of environmental stresses on fruit colour. While the term 'stress' usually conjures up negative connotations, moderate stress can enhance the concentration of numerous phytochemicals, including those that colour our fruits. The timing, duration and severity of any stress will determine the plant's response and consequent impact on colour.

Fruit pigmentation is part of the programmed development that culminates in ripening. This orchestrated process is largely driven by phytohormones and mediated via transcription factors (TFs) (Allan & Espley, 2018; Kumar et al., 2013). This process, however, can be strongly influenced by environmental conditions such as light intensity and spectral quality as well as temperature. Knowledge of

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Plant, Cell & Environment* published by John Wiley & Sons Ltd.



FIGURE 1 A range of pigments, including anthocyanins and carotenoids, confer numerous health benefits to the consumer. [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1111/pcel.14684)]

how the environment affects fruit colour will become increasingly important under changing climatic conditions globally. Increases in solar radiation and temperature as well as less predictable rainfall and a higher incidence of drought are predicted to affect crops (Challinor et al., 2014; Lobell & Gourdji, 2012). Climate change is expected to require the breeding of new varieties adapted to different conditions to tackle the potential issues of compromised plant health, flowering, fruit set and fruit development.

Colours in fruit not only protect the fruit itself, but also have implications for human health. As such, they have been recognised as an integral component of a healthy diet (Martin & Li, 2017). In fact, rather than simply maintain the amounts of these pigmented phytochemicals, many consider that they should be enhanced to promote greater potential health benefits (D'Amelia et al., 2018; Poiroux-Gonord et al., 2010).

2 | FRUIT PIGMENTS AND THEIR EFFECTS ON HUMAN HEALTH

One of the key drivers for research into plant pigments is the association of these compounds with dietary health (Martin & Li, 2017). Fruits are particularly rich sources of these phytochemicals and so changes in the composition or concentration are of interest. For carotenoids, one of the most abundant and important of these pigmented phytochemicals is β -carotene, which has provitamin A properties and is converted into retinol (vitamin A) in the body. Other carotenoid sources of provitamin A include α -carotene and β -cryptoxanthin (Rodriguez-Concepcion et al., 2018). Vitamin A is an essential part of our diet and deficiency is still a major problem in developing countries (West & Darnton-Hill, 2008), which prompted the development of provitamin A-enriched crops (Giuliano, 2017). Carotenoids also provide the macular pigments, xanthophylls including lutein and zeaxanthin, which have antioxidant and short-wavelength light absorption functions in the eye (Bernstein et al., 2016). These are associated with eye health, particularly the prevention of age-related macular degeneration (Carpentier et al., 2009). Consumption of carotenoids has also been associated with disease prevention, including a reduced incidence of

cardiovascular disease and some cancers, as well as positive effects on cognitive function (Leermakers et al., 2016; Lindbergh et al., 2018; Mein et al., 2008).

An increasing number of clinical trials have shown evidence for the health claims associated with anthocyanins. Many of these studies link anthocyanins to protection against human diseases, including cardiovascular and age-related degenerative diseases, obesity, metabolic syndrome and cancer (Gonçalves et al., 2021; Kalt et al., 2020; Pojer et al., 2013). There was earlier speculation about the mechanism of action, as these compounds were not found in human circulatory systems at the site of the action, but it appears now that bioavailability is greater than originally thought (Kay et al., 2017; Lila et al., 2016). If the protection that pigments provide in fruit is due to their antioxidant activities, is this also the mechanism the major human health benefits associated with their dietary consumption? This is less certain, and further research is required to confirm the modes of action. However, it has recently been established that one key mechanism for the health effects is via the extensive impact of an anthocyanin-rich diet on the gut microflora, which further process the final functional compounds (Espley et al., 2014; Groh et al., 2020; Jucá et al., 2020; Liso et al., 2018; Ntemiri et al., 2020; Ou et al., 2020).

3 | THE STRESS RESPONSE: A BRIEF OVERVIEW

When plants are subjected to sufficient environmental stress, reduction-oxidation (redox) reactions occur (Figure 2). Under normal conditions, partially reduced or activated forms of oxygen, known as reactive oxygen species (ROS), are generated, but at a rate that can be managed by the plant's ROS-scavenging machinery. Under highly stressful conditions, this redox homeostasis can become unbalanced, and these highly reactive molecules can cause cellular damage. As cellular ROS concentrations increase, for example with extreme temperature or drought, damage can afflict DNA, RNA, membranes and ultimately destroy the cell itself (Mittler, 2002). As oxygen is produced during photosynthesis, the chloroplastic photosynthetic machinery, particularly in leaves, is the site of much of the ensuing




FIGURE 2 The major sources of environmental stress that affect fruit pigmentation. (a) Examples of sources of environmental stress, (b) responses to stress as initiated by ROS production, and (c) examples of the stresses that can increase or decrease concentration of fruit pigments in apple, grape, tomato and mango. ROS, reactive oxygen species. [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1111/pce.14684)]

ROS (Foyer, 2018). In green developing fruit, where active photosynthesis is driving metabolism, this can still apply, but as the fruit ripens, ROS can be generated in other organelles, particularly mitochondria (Schertl & Braun, 2014). This rise in respiratory sourced ROS, may occur when the common chloroplast/chromoplast shift removes the central mechanisms for coping with ROS: the xanthophyll (and other carotenoid) cycles embedded within the photosynthetic machinery.

Since these redox reactions are inevitable in an oxygenated world, plants have had to evolve coping mechanisms. One mechanism is enzymatic, with enzymes such as superoxide dismutase (SOD), ascorbate peroxidase (APX), catalase (CAT), glutathione peroxidase, monohydroascorbate (MDHAR) and dehydroascorbate reductase (DHAR) (Das & Roychoudhury, 2014). The other major mechanism involves the non-enzymatic antioxidants such as ascorbic acid (AsA) and numerous polyphenols and terpenes (Decros et al., 2019). AsA is an important scavenger of reactive oxygen and in a scenario where stress leads to a reduction in photosynthesis and an increase in photooxidative damage and ROS production, AsA production may be enhanced. AsA production does not directly affect the colour of fruit and will not be covered in this review, but nevertheless it is interesting to consider that the same signals that mediate changes in the concentration of AsA in fruit, may also mediate changes in polyphenol and terpene production. The

polyphenols include flavonoids such as catechin, quercetin and anthocyanins, while terpenes include the pigmented carotenoids lycopene and β -carotene. It is clear then, that these plant pigments have a role in fruit beyond facilitating seed dispersal since there is good evidence for the induction of red pigmentation in vegetative tissues directly in response to ROS production. Plants have evolved this pigmentation to act as sunscreens and antioxidants and alleviate stresses from damaging environmental conditions (Davies et al., 2022).

The amount of ROS will also affect the concentrations of plant hormones, including abscisic acid (ABA), brassinosteroids (BRs), auxin and gibberellins (GAs). Phytohormones play a major role in ripening, with ethylene and ABA having key roles in climacteric and non-climacteric fruits, respectively (Fenn & Giovannoni, 2021). Both ethylene and ABA play dual roles in the fruit, being not only linked with ripening, but also with the stress response (Husain et al., 2020; Leng et al., 2013). Several studies have shown a connection between increased content of fruit ABA, as a response to stress or at onset of fruit ripening, and the accumulation of anthocyanins (Jia et al., 2011; Karppinen et al., 2018; Wheeler et al., 2009; Zifkin et al., 2012). It has been suggested that accumulation of ABA inside the cell induces downstream signalling involving ROS and calcium as secondary messengers, resulting in activation of the TFs controlling the anthocyanin biosynthetic pathway (Vighi et al., 2019).

Phytohormones can also interact, sometimes antagonistically such as has been shown for ABA and ethylene, but sometimes positively through providing an enhanced response to abiotic stresses (Müller, 2021).

In this review, we concentrate on the major environmental stressors that influence carotenoid and anthocyanin production in fruit crops. Chlorophyll accumulation and degradation also contribute to fruit colour, particularly before ripening occurs, but this falls outside the scope of this review. However, it should be recognised that before fruit ripening, the colour of fruit due to chlorophyll content may be affected by environmental stresses, such as heat and cold stress, for example as evident in tomato (Mesa et al., 2022). While betalains are also important plant pigments, they are most commonly found in leaves, flowers, tubers and only quite rarely found in fruit, such as in prickly pear (*Opuntia ficus-indica*) and dragon fruit/pitaya (*Hylocereus costaricensis*) (Cheek et al., 2022). They too are elevated in response to various environmental stresses: this has recently been well reviewed (Davies et al., 2018; Li et al., 2019) and will not be covered here.

4 | THE BIOSYNTHESIS AND TRANSCRIPTIONAL REGULATION OF FRUIT PIGMENTS

4.1 | The biosynthesis of flavonoid pigments

Anthocyanins are key pigments in fruit, generating red, blue and purple hues. These compounds usually accumulate in the epidermal or sub-epidermal tissues of different plant organs, offering protection against variable environmental stress factors, especially excess of light. The biosynthesis of anthocyanins occurs in the cytosol via the phenylpropanoid and flavonoid pathways. The precursors for flavonoids, including the pigmented anthocyanins, are malonyl-CoA and *p*-coumaroyl-CoA. From these precursors the first committed enzyme, chalcone synthase (CHS), forms chalcone, establishing the C₁₅ backbone (Figure 3). Chalcone is isomerised by chalcone isomerase (CHI) to produce chalcone naringenin, which is hydroxylated by flavanone 3 β -hydroxylase (F3H), converting naringenin to dihydroflavonol. Further hydroxylation is carried out by flavonoid 3'-hydroxylase




FIGURE 3 Simplified overview of the biosynthetic pathway leading to anthocyanin biosynthesis in fruit. Enzyme abbreviations as follows; PAL, phenylalanine ammonia lyase; C4H, cinnamate 4-hydroxylase; 4CL, 4-coumarate-CoA ligase; CHS, chalcone synthase; CHI, chalcone isomerase; F3H, flavanone 3-hydroxylase; F3'H, flavonoid 3' hydroxylase; F3'5'H, flavonoid 3',5' hydroxylase; FLS, flavonol synthase; OMT, O-methyltransferase; LAR, leucoanthocyanidin reductase; ANR, anthocyanidin reductase; DFR, dihydroflavonol 4-reductase; ANS, anthocyanidin synthase; UFGT, uridine diphosphate (UDP)-glucose:flavonoid 3-O-glucosyltransferase. [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com)]

(F3'H) to form cyanidin, or flavonoid 3'5'-hydroxylase (F3'5'H) to form delphinidin. Dihydroflavonol is reduced by dihydroflavonol 4-reductase (DFR) to produce leucoanthocyanidin, which is converted into the pigmented compound anthocyanidin by anthocyanidin synthase (ANS), also sometimes referred to as leucoanthocyanidin dioxygenase (LDOX). Finally, glycosylation is carried out by uridine diphosphate (UDP)-glycosyl:flavonoid 3-O-glycosyltransferase (UGT). Further addition of sugars and acyl side groups may follow. After biosynthesis, anthocyanins are transported to vacuoles or, rarely, cell walls. The differences in anthocyanin colours result from various factors, including the number of hydroxyl groups on the B-ring, the sugars and acyl side groups, the vacuolar pH or the accumulation of specific metal ions. To date, more than 650 different anthocyanins have been identified, the most common ones being glucosides of six aglycons: cyanidin, delphinidin, pelargonidin, petunidin, peonidin and malvidin (Jaakola, 2013; Zhang et al., 2014).

4.2 | The biosynthesis of carotenoid pigments

Carotenoids provide many of the red, orange and yellow pigments in fruit, as seen in citrus (*Citrus* spp.), tomato (*Solanum lycopersicon*), capsicum/pepper (*Capsicum* spp.) and papaya (*Carica papaya*). Carotenoids are essential for photoprotection and integral to the photosynthetic machinery and in green tissues are co-ordinately regulated with chlorophylls (Frank & Cogdell, 1996). The pathway is located in plastids and derived from the mevalonate-independent pathway (MEP) pathway, which generates the C₂₀ compound geranylgeranyl pyrophosphate (GGPP) (Cunningham & Gantt, 1998). The condensation of two GGPP molecules provides the first carotenoids, the colourless phytoene, and is catalysed by phytoene synthase (PSY) (Figure 4). This is often seen as the rate-limiting step for carotenoid synthesis, regulating downstream pathway flux. A two-step desaturation of phytoene, by phytoene desaturase (PDS), catalyses phytoene into phytofluene and then zeta-carotene, which is converted to di-cis zeta-carotene by zeta-carotene isomerase (Z-ISO). Desaturation by zeta-carotene desaturase (ZDS) and isomerization by carotene isomerase (CRTISO) produces lycopene and from this the pathway branches, with lycopene beta cyclases (LCYB) producing alpha-carotene and the nutritional vitamin A precursor, beta-carotene (Moise et al., 2014). The conversion of alpha-carotene to lutein and beta-carotene to zeaxanthin is through the action of carotene hydroxylases. Beyond these steps, other carotenoids such as xanthophylls, including astaxanthin, are generated and finally, through the actions of nine-cis-epoxycarotenoid dioxygenases (NCEDs) and carotenoid cleavage dioxygenases (CCDs), the phytohormone abscisic acid (ABA) and a range of apocarotenoids are produced (Giuliano, 2014). Carotenoid accumulation is further influenced by cell physiology, specifically the number of available carotenoid storage plastids (Cazzonelli & Pogson, 2010).




FIGURE 4 Simplified overview of the carotenoid biosynthetic pathway in fruit. Enzyme abbreviations as follows; PSY, phytoene synthase; PDS, phytoene desaturase; Z-ISO, ζ-carotene isomerase; ZDS, ζ-carotene desaturase; CRTISO, carotenoid isomerase; LCYE, lycopene ε-cyclase; LCYB, lycopene β-cyclase; BCH, β-carotene hydroxylase; CYP97, cytochrome P450-type monooxygenase 97; ZEP, zeaxanthin epoxidase; CCS, capsanthin-capsorubin synthase; NXS, neoxanthin synthase. [Color figure can be viewed at wileyonlinelibrary.com]

4.3 | The transcriptional regulation of plant pigments

The transcriptional regulation of anthocyanins in fruit has been well documented: at its core is a TF complex, termed the MYB-bHLH-WD40 (MBW) complex (Allan et al., 2008). The MYB component is derived from the subgroup (SG) 6 class of MYBs (as defined by Stracke et al., 2001) as opposed to the SG5 MYBs, which control proanthocyanidins and has been characterised in many fruit species (Albert et al., 2014; Jaakola, 2013). The MYBs are expressed in ripening fruit and activate either the whole biosynthetic pathway or a subset of genes in the pathway. Examples of activators include FaMYB10 and FvMYB10 in strawberry (Lin-Wang et al., 2014; Medina-Puche et al., 2014), MYB10 in nectarine (*Prunus persica*) (Ravaglia et al., 2013), VmMYBA in bilberry (*Vaccinium myrtillus*) (Karppinen et al., 2021), PpMYB10.1 in peach (*Prunus persica*) (Rahim et al., 2014), VcMYBA1 in blueberry (*Vaccinium Section Cyanococcus*)

(Plunkett et al., 2018) and MYB114 in pear (*Pyrus bretschneideri*) (Yao et al., 2017). MYBs can also act as repressors of anthocyanin, such as the SG4 MYBs, FaMYB1 from strawberry (Aharoni et al., 2001) and MdMYB16 from apple (*Malus domestica*) (Xu et al., 2017), as recently reviewed by (LaFountain & Yuan, 2021). There is increasing evidence for the interactions of different types of MYBs on anthocyanin regulation, such as the co-ordinated actions of MYBA1 and MYBPA1.1 in blueberry (Lafferty et al., 2021). Transcriptional regulation of anthocyanin is not restricted to the MBW complex: other TFs, such as MADS VmTRD4 in bilberry (Jaakola & Poole, Jones, et al., 2010) and PpNAC1 in peach (Zhou et al., 2015) have been shown to have regulatory control. This TF-mediated control of pigments occurs in response to both developmental cues, but also environmental stresses.

Carotenoid biosynthesis is stimulated by the presence of ROS (Pan et al., 2009). Stress perceived in the leaves can trigger responses in the fruit, for example, the incidence of high light in leaves of citrus leads to increases in carotenoids in the fruit (Poiroux-Gonord et al., 2013). Many TFs have been associated with carotenoid biosynthesis as part of the fruit ripening cascade, but the evidence for there being master transcriptional regulators of carotenoid pigments is less clear cut than for anthocyanins (Stanley & Yuan, 2019). A number of TFs have recently been shown to control at least parts of the pathway. In citrus, the AGAMOUS-LIKE MADS TF, CsMADS6, was shown to activate the promoter of *LCYb1* and drive the expression of carotenoid pathway genes *PSY*, *PDS* and *CCD1*, while repressing transcription of *LCYE* and so increasing the flux towards β -carotene (Lu et al., 2018). Another TF in citrus, CrMYB68, negatively regulates the expression of beta-carotene hydroxylase (BCH), controlling the α - and β -branches of carotenoid biosynthesis (Zhu et al., 2017). A MYB from kiwifruit (*Actinidia chinensis* var. *deliciosa*), MYB7, was reported to activate the promoter of *LCYb* and in heterologous systems elevated expression of *PSY*, *PDS* and *ZDS* to drive carotenoid accumulation (Ampomah-Dwamena et al., 2019). In papaya, two other TF classes have been implicated in regulating carotenoids: CpbHLH1 and CpbHLH2 repress and activate respectively the promoters of *CYCB* and *LCYB*, while the NAC TFs, CpNAC1 and CpNAC2, bind the promoters of *PDS* (NAC1) and *PDS*, *ZDS*, *LYCE* and *BCH* (NAC2) (Fu et al., 2016; Fu et al., 2017). The greatest body of work on the transcriptional control of carotenoids has been carried out in tomato where the major ripening MADS TFs, Tomato AGAMOUS-LIKE (TAGL1, homologous to CsMADS6), Ripening Inhibitor (RIN), FRUITFULL1 (FUL1) and FUL2 exert largely positive effects on the carotenoid pathway, as reviewed in Stanley & Yuan (2019).

The co-ordinated fruit ripening process involves metabolic or phytohormonal signals that drive both the carotenoid and anthocyanin pathways. There are ways in which these pathways are intrinsically linked, such as the carotenoid-derived phytohormone ABA, which can induce anthocyanin production. In this review, we highlight some stresses that do result in changes to the concentration of both anthocyanins and carotenoids, suggesting a common signal. An intriguing question remains as to whether a master transcriptional

regulator exists. Interestingly the two pathways have been shown to respond to one gene, the negative regulator of photomorphogenic response, DET1 (De-etiolated 1). When DET1 was suppressed by RNAi in tomato fruit, there was an increase in both carotenoids (lycopene and β -carotene) and flavonoids (chlorogenic acid, naringenin-chalcone and quercetin) (Davuluri et al., 2005). Whilst this did not affect anthocyanin, it showed that DET1 influenced both pathways simultaneously.

4.4 | Influence of gene \times environment

Whilst the general biosynthesis and regulation of colour pathways has been elucidated, the final colour of the fruit is determined by both genetic and environmental factors. The genetic background determines the ability of fruit to produce certain pigments, whilst environmental factors affect the final composition and amounts of pigments in a ripe fruit. The fruit skin, which acts as a barrier protecting the internal tissues of developing fruit, including the photosynthetic machinery, is where the greatest effects of environmental factors on fruit pigment production are normally detected. Both controlled and field studies with cultivated and wild ecotypes have provided a better understanding of gene \times environment (G \times E) interaction in fruit pigment production (Jaakola & Hohtola, 2010). Such studies have identified variation in anthocyanin content between fruit ecotypes or cultivars growing in different geographical locations (Fredes et al., 2014; González-de-Peredo et al., 2019; Sikuten et al., 2021). With wild *Vaccinium* berries, field studies have often shown high variation in the anthocyanin composition between growth location and seasons (Connor et al., 2002; Rohloff et al., 2015). However, specific geographical trends also have been observed, such as higher total anthocyanin contents and a higher proportion of delphinidin branch anthocyanins towards northern latitudes (Åkerström et al., 2010; Lätti et al., 2008, 2010) and at higher altitudes (Zoratti et al., 2015). A controlled study with southern and northern bilberry ecotypes indicated that ecotypes from higher altitudes produced a higher concentration of anthocyanins irrespective of day length and temperature conditions (Uleberg et al., 2012). Geographical variation has also been detected in the carotenoid contents of different fruit species and cultivars (Dias et al., 2021). In some cases, climatic factors have been found to have greater impacts on carotenoid biosynthesis than cultivar differences, for instance in papaya (Kimura et al., 1991), mango (*Mangifera indica*) (Mercadante & Rodriguez-Amaya, 1998), and West Indian cherry (*Malpighia glabra*) (Rodriguez-Amaya, 2001). A comprehensive analysis of the carotenoid content in Portuguese fruit showed variation between the same apple and orange (*Citrus sinensis*) cultivars grown in different geographical regions (Dias et al., 2009). High variation in carotenoid content was detected between wild Seabuckthorn (*Hippophae rhamnoides*) populations growing at different altitudes trans-Himalaya in India (Korekar et al., 2014). The major stressors are considered in the following sections, with the proviso that G \times E can have additional and compounding effects.

5 | THE MAJOR ENVIRONMENTAL STRESS FACTORS THAT AFFECT FRUIT COLOUR

5.1 | Light stress

Light is one of the most important environmental factors affecting the accumulation of pigments during fruit ripening. Light effects can be categorised by duration (photoperiod), intensity (quantity), and quality (wavelength), all of which have been shown to affect pigment production. Light stress refers to the detrimental effect that excess or inadequate light intensities and their fluctuations can have on plant function and development. Excess of light can damage the photosynthetic machinery through the increased ROS concentrations, whereas insufficient light limits the photosynthetic activity. In addition, fluctuations in light wavelengths, especially excess of UV-light, can also cause light stress. Stress from non-optimal light conditions can cause alterations in pigment production (Zoratti et al., 2014). Plants convert photons of light through photosynthetic fixation of carbon dioxide into sugars. Light quality affects both photosynthetic rate and assimilation. Exquisite mechanisms have been developed to tolerate light stress, accumulation of pigments being one of the most important (Yang et al., 2019).

Little or moderate light stress can actually be beneficial for fruit quality by enhancing pigment concentration (Table 1). On-tree apple experiments have shown that light stress induces anthocyanin production: these anthocyanins may reduce the amount of excess light as well as reducing the damage from the increased ROS produced to protect chlorophylls and carotenoids as part of the photooxidative machinery (Merzlyak & Chivkunova, 2000). On-tree experiments have also shown that full spectra, including UV light, are essential for anthocyanin production (Henry-Kirk et al., 2018). Fruit bagging and shading experiments confirm the importance of direct exposure to light for biosynthesis of anthocyanins and carotenoids. Light exclusion results in transcriptional suppression of biosynthetic pathway genes and reduced accumulation of pigments (Downey et al., 2006; Saini & Keum, 2018; Zhu et al., 2021; Zoratti et al., 2014). Whereas many fruits, like apples, have an absolute light requirement for anthocyanin biosynthesis (An et al., 2020a), there are species and cultivars that show no change in anthocyanin accumulation when in shaded conditions or any degradation of anthocyanins under high irradiation (Downey et al., 2006; Zoratti et al., 2014). This indicates that there are different genetic regulation mechanisms for anthocyanin biosynthesis in some fruits, which are not fully understood.

Anthocyanins intercept and absorb light energy to avoid photo-damage caused by excess light energy, directly scavenge free radicals and indirectly remove ROS by interacting with molecules in other signalling pathways (Li et al., 2019). In extreme high light conditions, anthocyanins play an important role in mitigating photoinhibition and photodamage. Naturally red-skinned pear species have been shown to maintain better photosynthetic capacity under high light and high temperature conditions, while some green-skinned pear cultivars

increase skin anthocyanins in response to high light stress (Thomson et al., 2018).

In tomato, the accumulation of carotenoids during ripening is significantly influenced by light. Phytochrome induces degradation of phytochrome-interacting factor 1 (PIF1), a bHLH-type TF. In mature green tomato, PIF1 directly binds to the PIF-binding E-box PBE box of the PSY1 promoter of the carotenoid biosynthetic pathway, to repress carotenogenesis. During ripening, developmentally controlled degradation of chlorophylls reduces the self-shading effect, allowing phytochrome-mediated degradation of PIF1, which leads to depression of PSY1. This accelerates carotenoid biosynthesis and shifts the profile from xanthophyll, which is typical for leaves, to carotenes, mainly lycopene and β -carotene (Saini & Keum, 2018). Exposure of tomato to excessive sunlight has been reported to inhibit the synthesis of lycopene (Brandt et al., 2006; Ilić & Fallik, 2017). It has been proposed that ROS and redox status, combined with sugar and carbon status, integrate the stress response and photosynthetic metabolism, and so influencing the synthesis of carotenoids (Fanciullino et al., 2014). In Clementine (*Citrus clementina*), accumulation of high amounts of soluble sugars through photosynthesis during the early stages of fruit development negatively influenced plastid development, which affected carotenoid accumulation, although later in fruit development, greater sugar accumulation increased carotenoid production (Fanciullino et al., 2014; Saini & Keum, 2018).

The quality of the light spectrum has been shown to affect fruit pigments. Treatments with UV-B and UV-C light during ripening or postharvest resulted in increased anthocyanin concentrations in many fruit species, including apple, peach, wine grape (*Vitis* spp.), and blueberry (Chen et al., 2021; Henry-Kirk et al., 2018; Ubi et al., 2006; Zhao et al., 2017; Zoratti et al., 2014). Changes in carotenoid content following exposure to UV-light have also been reported in tomato (Giuntini et al., 2005) and wine grape berries (*Vitis vinifera*) (Joubert et al., 2016).

Solar UV-B radiation has extensive photobiological effects on plants: stress caused by UV-B is known to enhance the production of ROS, leading to damage of DNA, proteins and the photosynthetic apparatus. In plants, UV-B light is absorbed by tryptophan amino acid residues in the dimeric form of UVB photoreceptor, leading to UVR8 monomerization and formation of a complex with CONSTITUTIVE PHOTOMORPHOGENIC1 (COP1), a known mediator of diverse light signals in plants interacting with regulators of both anthocyanin and carotenoid biosynthesis (Stanley & Yuan, 2019; Zoratti et al., 2014). Plant pigments have an important role in protection of tissues against UV-radiation. For instance, anthocyanins and flavonols have been shown to accumulate in the epidermal tissues, such as fruit skin, to shield against excess UV-light (Zoratti et al., 2014). During photosynthesis, carotenoids harvest light energy and transfer this energy to chlorophylls through singlet-singlet excitation transfer, a lower energy state transfer. Excessive energy from chlorophylls is absorbed by carotenoids through triplet-triplet transfer, which is a higher energy state transfer essential for photo-protection. Carotenoids with more than eleven conjugated double bonds have a high capacity

TABLE 1 Examples of the effect of major environmental stresses (light, temperature and drought) on the concentration of pigments in different species of fruit.

Stress	Species	Fruit pigment	References
UV-B, UV-C	Apple (<i>Malus domestica</i>)	Anthocyanin ↑ (Cyanidin)	Fang et al. (2019); Henry-Kirk et al. (2018); Ubi et al. (2006)
	Tomato (<i>Solanum peruvianum</i>)	Anthocyanin ↑	Tapia et al. (2022)
	Peach (<i>Prunus persica</i>)	Anthocyanin ↑	Zhao et al. (2017)
High light	Apple (<i>M. domestica</i>)	Anthocyanin ↑ carotenoids ↑	Merzlyak & Chivkunova (2000)
Red/Blue light	Strawberry (<i>Fragaria ananassa</i>)	Anthocyanin ↑	Miao et al. (2016)
	Bilberry (<i>Vaccinium myrtillus</i>)	Anthocyanin ↑	Samkumar et al. (2021)
	Chili pepper (<i>Capsicum annuum</i>)	Carotenoids ↑ (β-carotene)	Pola et al. (2019)
Red light	Capsicum (<i>C. annuum</i>)	Carotenoids ↑	Lopez et al. (1986)
	Kumquat (<i>Fortunella crassifolia</i>)	Anthocyanin ↑	Gong et al. (2021)
Blue light	Capsicum (<i>C. annuum</i>)	Carotenoids ↓	Lopez et al. (1986)
Heat	Apple (<i>M. domestica</i>)	Anthocyanin ↓ (Cyanidin)	Lin-Wang et al. (2011)
	<i>Malus profusion</i>	Anthocyanin ↓ (Cyanidin)	Rehman et al. (2017)
	Grape (<i>Vitis vinifera</i> L.)	Anthocyanin ↓ (Malvidin, petunidin, cyanidin, peonidin)	Movahed et al. (2016)
	Grape (<i>Vitis lambrusca</i> x <i>Vitis vinifera</i>)	Anthocyanin ↓ (Total)	Yamane et al. (2006)
	Grape (<i>Vitis lambruscana</i>)	Anthocyanin ↓ (Malvidin, delphinidin, petunidin, peonidin)	Ryu et al. (2020)
	Tomato (<i>Lycopersicon esculentum</i>)	Lycopene ↓	Brandt et al. (2006); Tomes (1963)
Cold	Apple (<i>M. domestica</i>)	Anthocyanin ↑ (Cyanidin)	Chen et al. (2021); Fang et al. (2019); Ubi et al. (2006)
	Black currants (<i>Ribes nigrum</i>)	Anthocyanin ↑	Woznicki et al. (2015)
	Cloudberries (<i>Rubus chamaemorus</i>)	Anthocyanin ↑	Martinussen et al. (2010)
	Blood orange (<i>Citrus sinensis</i>)	Anthocyanin ↑	Butelli et al. (2012); Sicilia et al. (2020)
	Mango (<i>Mangifera Indica</i> L)	Carotenoids ↓ (violaxanthin, β-carotene)	Rosalie et al. (2018)
Drought	Apple (<i>M. domestica</i>)	Anthocyanin ↑	An et al. (2020c)
	Grape (<i>V. vinifera</i> L.)	Anthocyanin ↑	Bucchetti et al. (2011); Castellarin et al. (2007); Deluc et al. (2009); Santesteban et al. (2011)
	Strawberry (<i>F. ananassa</i>)	Anthocyanin ↑	Perin et al. (2019)
	Tomato (<i>Solanum peruvianum</i>)	Anthocyanin ↑	Tapia et al. (2022)
	Jujube (<i>Ziziphus jujuba</i>)	Anthocyanin ↓ Carotenoids ↓	Jiang et al. (2020)
	Tomato (<i>L. esculentum</i>)	Carotenoids ↓	De et al. (2007)
	Tomato (<i>L. esculentum</i>)	Carotenoids ↑ (Lycopene, β-carotene)	Favati et al. (2009); Klunklin & Savage (2017)

Note: Arrows represent increase ↑ or decrease ↓ in total pigments (individual compounds when information available).

for quenching singlet oxygen produced under light stress (Maoka, 2020).

Other light spectrum wavelengths have also been shown to influence the accumulation of fruit pigments and particularly red and blue wavelengths affect the biosynthesis of anthocyanins and

carotenoids. Red or blue light treatment has resulted in a detectable increase in anthocyanin content in several fruit crops, including strawberries, grape, pear, apple, and cherry (*Prunus avium*) (Kokalj et al., 2019a; 2019b; Koyama et al., 2012; Miao et al., 2016; Tao et al., 2018). Recently, both red and blue light were shown to increase

anthocyanin biosynthesis during bilberry ripening (Samkumar et al., 2021, 2022). Interestingly, differences in the perception of red and blue light were detected between berries ripening whilst remaining attached to the mother plant, or when detached in Petri dishes. Transcriptomic analysis revealed differences in red and blue light signalling, leading to altered anthocyanin content in the ripe, treated berries, with red light increasing the anthocyanin content 12-fold compared with the control (Samkumar et al., 2021).

Many studies have shown that red light typically increases carotenoid biosynthesis in fruits, while blue light has less effect, although there are differences between species, cultivars and ripening stages. Spectral light quality experiments monitoring carotenoid biosynthesis in ripening *Capsicum annuum* showed that red radiation was the most effective in increasing carotenoid accumulation, whereas green and blue radiation inhibited the formation of capsanthin, the major carotenoid influencing fruit colour (Lopez et al., 1986; Pola et al., 2019). Similar results have been shown in *Citrus* (Gong et al., 2021; Ma et al., 2012). In kumquat (*Fortunella crassifolia*) fruit, a NAC family TF, FrCNAC22, was shown to mediate red light-induced carotenoid accumulation (Gong et al., 2021). Moreover, PIF1 regulation, following blue light perception, has been shown to act as a negative regulator of carotenoid biosynthesis (Gong et al., 2021).

5.2 | Temperature stress

Temperature stress in plants is classified in three types that may be too high, chilling or freezing temperatures all affecting the ROS machinery in the plant cells. Plants respond to temperature stress by regulating membrane lipid composition, by expressing stress related TFs, and by detoxification pathways and metabolite biosynthesis (Kai & Iba, 2014). Both below and above optimal growth temperatures, have been shown to affect plant pigment accumulation during fruit ripening. Although there is broad variation in what is the optimal temperature depending on species, cultivar and ecotype, heat stress can inhibit pigment biosynthesis in ripening fruits. Studies demonstrate that this is the case for both carotenoid and anthocyanin pigments (Gashu et al., 2022; Kliewer, 1977; Lin-Wang et al., 2011; Tomes, 1963; Ubi et al., 2006) (Table 1). When heat stress is moderate, the changes can be reversible, whereas severe episodes can cause irreversible changes to metabolites, resulting in crop failure.

Heat stress has been shown to inhibit anthocyanin biosynthesis in grape berry skin in controlled studies with temperatures higher than 30°C (Ryu et al., 2020; Yamane et al., 2006). Active inhibition and degradation of anthocyanin biosynthesis under high temperature conditions have also been reported in other fruits such as *Malus profusion* (Rehman et al., 2017) as well as in grape (Movahed et al., 2016). Changes in ABA contents and in ABA/GA ratio have been associated with changes in anthocyanin concentrations under different temperature conditions (Ryu et al., 2020). Hot conditions during the growing season can dramatically reduce anthocyanin

production in numerous apple cultivars, with an associated reduction in the transcription of the biosynthesis genes (Ubi et al., 2006) and the pathway regulating MYB TF, MYB10 (Lin-Wang et al., 2011). This appeared to be repression of anthocyanin activation, rather than an increase in transcription of known anthocyanin MYB repressors.

Almeida et al. (2021) found that ripe tomato fruit had better plasticity to restore carotenoids following a heat wave compared with tomatoes that experienced excessive heat during the breaker stage (when pink colour is just starting to develop). They concluded that heat-related transcriptional and posttranscriptional misregulation during early carotenogenesis affected the carotenoid composition in the ripe tomatoes. Moreover, upstream ripening regulators, such as CNR and NOR, were also affected by the heat treatments. The results also suggested that epigenetic mechanisms were mediating heat-induced transcriptional changes.

It is not only high temperatures that can be problematic for fruit colour. Low temperature conditions at around 4–10°C can cause stress in developing fruits because of ROS production and generation of malondialdehyde (MDA) in the cell, which can lead to severe membrane lipid peroxidation. Cold stress also promotes the accumulation of anthocyanin- and carotenoid-regulating phytohormones, ABA, jasmonates (JA), ethylene (ETH) and salicylic acid (SA) (He et al., 2022). ABA, JA and polyamines have been reported to be associated with low-temperature stress tolerance in apple fruitlets (Yoshikawa et al., 2007).

Lower temperatures have been observed to increase anthocyanin biosynthesis in many fruits, such as apples, table grapes, black currants (*Ribes nigrum*), cloudberries (*Rubus chamaemorus*), while anthocyanin accumulation is inhibited during heat stress (Chen et al., 2021; Downey et al., 2006; Martinussen et al., 2010; Woznicki et al., 2015). In blood orange (*Citrus sinensis*), anthocyanin production has been shown to be dependent on cold (Butelli et al., 2012; Sicilia et al., 2020). An increase in anthocyanin concentration in fruit skin provides some tolerance to cold stress, for instance in mango (Sivankalyani et al., 2016; Sudheeran et al., 2018). This production of anthocyanin at lower temperatures provides an alternative means to other ROS coping mechanisms which are compromised by cold. However, for other fruits such as strawberry, temperatures below 10°C have been shown to inhibit anthocyanin biosynthesis, a process mediated by MITOGEN-ACTIVATED PROTEIN KINASE₃ (FvMAPK₃) (Mao et al., 2022).

Whereas high temperature clearly represses transcription of MYB10 in apple fruit, the opposite is seen with low temperature in crabapple. In this case, the transcription of the MBW activation complex, including MYB10, bHLH3/33 and TTG1, is elevated, leading to increased anthocyanin production in leaves (Tian et al., 2015). Other TFs are involved in this response and Fang et al. (2019) showed that the zinc finger TF, MdbBX20, responds to low temperature via MdbHLH3 interaction, to drive anthocyanin production in a HY5-dependent manner.

Cold stress has been shown to affect carotenoid contents in fruits as a response to the production of ROS, in a species-specific manner. For example, higher lycopene contents have been shown to

offer tolerance against cold stress in grapefruit (*Citrus paradisi*) (Lado et al., 2015), whereas in mango, carotenoid contents decreased under cold stress (Rosalie et al., 2018). In pepper (*C. annuum*) fruit, Zhang et al. (2020) showed that *CaATHB-12*, a member of the HD-ZIP I gene family, is one of the key regulators of fruit carotenoid content.

Diurnal temperature, with cool nights and warm days, is critical to optimal fruit colour in some species and a reduction in diurnal temperature range, as may become more evident with climate change, will impact anthocyanin in fruit peel, such as in apple (Qu & Zhou, 2016). Clearly, optimal day and night temperatures help to produce high quality coloured fruit, while temperatures outside of this range often positively (cold) or negatively (excess heat) effect the concentration of colour.

5.3 | Water stress

Water stress in plants may arise either due to excess of water or due to water deficit, which is the more common water stress, also known as drought-stress. Drought induced water stress leads to decreased water potential, downregulation of photosynthesis-related genes and reduced availability of CO₂ being also one of the major factors in the stress caused by excess of light (Osakabe et al., 2014). Water shortages are already a serious problem for much of the world's agriculture and horticulture (Lobell & Gourdji, 2012). The predicted increase in severe climate events due to climate change, including drought, is likely to present even greater challenges for growers. Of particular concern is the potential impact on global food supply chains of reduced yield of major grain crops (Araus et al., 2002). Predicting the outcomes of different water deficiency scenarios and how these will affect crops is complex (Tramblay et al., 2020). For essential photosynthetic pigments in leaves (carotenoids and chlorophylls), water deficiency can result in reduced growth and yield (Jaleel et al., 2009) and in some fruits, water deficiency can reduce both anthocyanin and carotenoid concentrations, reducing fruit colour (Jiang et al., 2020). However, for other fruits, water deficiency outcomes are not necessarily deleterious (Table 1). This may seem counter-intuitive since drought is likely to decrease leaf photosynthesis and, therefore, the distribution of primary metabolites to the fruit. These primary metabolites provide the precursors for all secondary/specialised metabolites and so, logically, may reduce the pool of substrates for pigmented compounds. However, since both the carotenoid and phenylpropanoid pathways are partly regulated by stress-induced ROS production, there are also likely to be positive effects of water deficiency on fruit pigments. The outcomes can be species-dependent and, given this tension between pathways, highly variable owing to the timing of fruit development at which the water deficiency occurs (cell division, cell expansion or ripening phases), and the severity and duration of the deficit. In terms of overall fruit quality, there are possible effects on other major sensory attributes such as aromas, the sugar/acid balance and texture (Ripoll et al., 2014).

In some cropping situations, water deficiency is intentionally used to improve colour. In red grape, anthocyanin is an important

determinant of wine quality. When water is deficient, grape berry size is reduced, creating a higher skin:flesh ratio which, when coupled with increased anthocyanin production, can improve wine quality (Gambetta et al., 2020). Increases in grape berry anthocyanin concentrations following exposure to water deficit have been reported in Cabernet Sauvignon (Deluc et al., 2009), Merlot (Bucchetti et al., 2011) and Tempranillo (Santesteban et al., 2011). Anthocyanin composition can also change in a cultivar-specific manner, such as the drought-induced increase in acylated anthocyanins in Cabernet Sauvignon, while the same conditions led to a reduction in these compounds in Syrah (Hochberg et al., 2015). Further compositional changes can be created by a shift towards more tri-hydroxylated anthocyanins (darker purples and blues) via the upregulation of flavonoid 3'5' hydroxylases (Castellarin et al., 2007).

Phytohormones are intrinsically linked with drought perception and the resultant transcriptional cascade (Ullah et al., 2018), particularly ABA (Yamaguchi-Shinozaki & Shinozaki, 2005). For example, in strawberry fruit, drought conditions increased ABA, which was correlated with elevated anthocyanin concentration (as well as AsA) without a reduction in fruit yield (Perin et al., 2019). A previous study in strawberry clearly demonstrated how the presence of ABA regulated the expression of the anthocyanin-regulating MYB TF, *FaMYB10*, which, in turn, elevated the anthocyanin biosynthetic genes and fruit colour (Medina-Puche et al., 2013). In apple, an alternative molecular model of drought-induced anthocyanin production has been proposed, whereby the ethylene response factor ERF38 partners with the homologous MYB1 to drive the anthocyanin biosynthetic pathway genes (An et al., 2020c). In wild tomato (*Solanum peruvianum*), water deficit results in a greater anthocyanin concentration in the fruit skin, partly mediated by the enhanced expression of MYB TF, *SpAN2* (Tapia et al., 2022).

For fruit colours derived from carotenoids the picture is less clear cut than for anthocyanins. In some cases, water deficit has been shown to reduce fruit carotenoid concentrations (De et al., 2007; Jiang et al., 2020). However, in tomato, there is strong evidence that concentrations of lycopene (and β -carotene) increase with water deficiency (Favati et al., 2009; Klunklin & Savage, 2017; Patané et al., 2021; Zushi & Matsuzoe, 1998).

5.4 | Nutrient stress

While fruit pigments are highly influenced by temperature, light and water, it is important to consider that they may also be altered by other environmental factors. Nutrient stress in plants can be caused by low or excess levels of minerals. Unbalanced nutrient supply may hinder the plant growth and performance, and an excess or low level of a particular mineral can affect the uptake of other nutrients. For instance, salinity stress can either increase or reduce tomato anthocyanins depending on cultivar, while the pigmented carotenoids are reliably increased by as much as 2–3-fold across a range of cultivars (Borghesi et al., 2011). However, perhaps the most important consideration is nitrogen (N) status (Wang et al., 2018).

In *Arabidopsis*, restricted N availability has been shown to shift metabolism towards flavonoid production, including anthocyanins, with evidence for increased transcription of the regulating TF (AtPAP1) and the biosynthetic pathway (Lillo et al., 2008). If this is also true for fruits, it suggests that horticultural practices can elevate fruit anthocyanin concentrations, but that in planning nutrient applications, a balance is required to avoid detrimental yield effects on the overall crop (Jezek et al., 2018). That the effects of N deficiency is not always so clear and in other crops, such as strawberry, and N application timing needs to be determined on a crop-by-crop basis (Jezek et al., 2018).

At the molecular level, studies in apple show a number of candidate genes linking N deficiency to an elevation in the anthocyanin pathway. Low N induces anthocyanin production, partly mediated by the mitogen-activated protein kinase (MAPK) signalling cascade, as shown for red-fleshed apples, where a MAPK family gene, *MdMKK9*, regulates the N status (Sun et al., 2022). Under low N conditions the autophagy nutrient recycling process can be activated, and studies in apple have demonstrated that the AuTophagy (ATG) protein, *MdATG18a*, improves tolerance to nitrogen deficiency via upregulation of the anthocyanin pathway (Sun et al., 2018). In these cases, the increases in anthocyanin content are driven by an upregulation of the anthocyanin-regulating TF, *MdMYB1/10*. Conversely, the BTB/TAZ protein, *BTB2*, which is lowly expressed under low N but highly expressed under high N, negatively regulates the MYB, suppressing anthocyanin accumulation (An et al., 2020b). It seems that fruit carotenoid contents, as shown in tomato, are either decreased or less affected by low nitrogen supply depending on the timing of the nitrogen supply reduction (Bénard et al., 2009; Hernández et al., 2020).

6 | CONCLUSIONS AND PERSPECTIVES

From research reviewed here, it is clear that some environmental stresses can enhance fruit colour: low temperature, high light (and UV) and a degree of water deficit can all be beneficial. Against this, the severity and timing of these stresses are critical in determining whether outcomes are positive or negative. Some of these stresses can be controlled horticulturally, such as limiting water supply, but climate change is outside grower control and thus likely to disrupt our current growing practices. Consumer desire for ever more healthy fruits is unlikely to dissipate—the pigments in fruits provide many of these health benefits, packaged with one of the strongest motivators for purchase—the attractiveness of colourful fruit. Of course, fruit cultivars, no matter how colourful, will also have to reach other key consumer expectations, such as texture, storability and flavour. As climate change creates less predictable temperatures across many fruit-growing regions, understanding how fruit pigmentation will be affected will become increasingly important if we are to maintain or improve fruit colour. For some fruit crops, indoor production may become a more attractive (and commercial) proposition to avoid crop losses due to damaging climatic events.

Various options are available to maintain fruit colour in this changing environment, including agronomic approaches (Poiroux-Gonord et al., 2010). Similarly, postharvest treatments can provide the potential to preserve and enhance fruit pigments (de Pascual-Teresa & Sanchez-Ballesta, 2008). In the future, more substantive advances may be required to augment the incremental changes that can be made by in-field or postharvest management. In some circumstances, fruit breeding programmes have reduced the genetic and phenotypic diversity in breeding populations. To address production challenges associated with severe climate change, incorporating new genetic diversity is likely to become essential to maintain or improve fruit crops: the use of crop wild relatives that are better adapted to environmental stresses should be considered to introduce beneficial traits. Ideally, this would be in conjunction with deep bioinformatic resources (high quality genomes, transcriptomes and metabolomes), and an emphasis on mapping and molecular markers to help to target 'resilience' traits and rapidly incorporate these in new cultivars that are better adapted to our changing environment. Breeding varies markedly between species and varieties, and there will be an increasing need for developing crops suited to (and tested at) different geographic and environmental locations. The complexity of GxE interactions on fruit pigmentation is complex, and more research into this area is required to understand the plant response and potential breeding strategies for fruit colour improvement.

'Traditional' genetic engineering has provided many examples of successfully increasing plant pigments (Martin & Li, 2017). Sophisticated molecular engineering of fruit can provide platforms for elevated amounts of specific phytochemicals, including pigments (Li et al., 2018). The new breeding technologies such as CRISPR provide untapped potential for modifying our fruit crops both to improve their nutritional benefit (colour) and to tackle changing growing conditions (Wang et al., 2019). It is likely that adoption of new breeding approaches will be required to maintain and enhance fruit pigments as we navigate environmental changes to fruit-growing regions across the globe.

ACKNOWLEDGEMENTS

We apologise to those authors whose work was unable to be included in this review owing to space limitations. Research in the laboratory of R.V.E. is supported by The New Zealand Government Ministry of Business, Innovation and Employment's Endeavour Fund, contract C11X1704 'Filling the Void' and for L.J. by NordPlant (NordForsk grant no. 84597). We thank our Plant & Food Research colleagues Kevin Davies for helpful advice on the manuscript and Elysia Arnold and Ben Lawrence for illustrations. Open access publishing facilitated by New Zealand Institute for Plant and Food Research Ltd, as part of the Wiley - New Zealand Institute for Plant and Food Research Ltd agreement via the Council of Australian University Librarians.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

ORCID

Richard V. Espley  <http://orcid.org/0000-0002-1732-3688>

Laura Jaakola  <http://orcid.org/0000-0001-9379-0862>

REFERENCES

- Aharoni, A., De Vos, C.H.R., Wein, M., Sun, Z., Greco, R., Kroon, A. et al. (2001) The strawberry FaMYB1 transcription factor suppresses anthocyanin and flavonol accumulation in transgenic tobacco. *The Plant Journal*, 28, 319–332.
- Åkerström, A., Jaakola, L., Bång, U. & Jäderlund, A. (2010) Effects of latitude-related factors and geographical origin on anthocyanidin concentrations in fruits of *Vaccinium myrtillus* L. (bilberries). *Journal of Agricultural and Food Chemistry*, 58, 11939–11945.
- Albert, N.W., Davies, K.M., Lewis, D.H., Zhang, H., Montefiori, M., Brendolise, C. et al. (2014) A conserved network of transcriptional activators and repressors regulates anthocyanin pigmentation in eudicots. *The Plant Cell*, 26, 962–980.
- Allan, A.C. & Espley, R.V. (2018) MYBs drive novel consumer traits in fruits and vegetables. *Trends in Plant Science*, 23, 693–705.
- Allan, A.C., Hellens, R.P. & Laing, W.A. (2008) MYB transcription factors that colour our fruit. *Trends in Plant Science*, 13, 99–102.
- Almeida, J., Perez-Fons, L. & Fraser, P.D. (2021) A transcriptomic, metabolomic and cellular approach to the physiological adaptation of tomato fruit to high temperature. *Plant, Cell & Environment*, 44, 2211–2229.
- Ampomah-Dwamena, C., Thrimawithana, A.H., Dejnopratt, S., Lewis, D., Espley, R.V. & Allan, A.C. (2019) A kiwifruit (*Actinidia deliciosa*) R2R3-MYB transcription factor modulates chlorophyll and carotenoid accumulation. *New Phytologist*, 221, 309–325.
- An, J.-P., Liu, Y.-J., Zhang, X.-W., Bi, S.-Q., Wang, X.-F., You, C.-X. et al. (2020a) Dynamic regulation of anthocyanin biosynthesis at different light intensities by the BT2-TCP46-MYB1 module in apple. *Journal of Experimental Botany*, 71, 3094–3109.
- An, J.P., Wang, X.F. & Hao, Y.J. (2020b) BTB/TAZ protein MdBt2 integrates multiple hormonal and environmental signals to regulate anthocyanin biosynthesis in apple. *Journal of Integrative Plant Biology*, 62, 1643–1646.
- An, J.P., Zhang, X.W., Bi, S.Q., You, C.X., Wang, X.F. & Hao, Y.J. (2020c) The ERF transcription factor MdERF38 promotes drought stress-induced anthocyanin biosynthesis in apple. *The Plant Journal*, 101, 573–589.
- Araus, J.L. (2002) Plant breeding and drought in C3 cereals: what should we breed for? *Annals of Botany*, 89, 925–940.
- Bénard, C., Gautier, H., Bourgaud, F., Grasselly, D., Navez, B., Caris-Veyrat, C. et al. (2009) Effects of low nitrogen supply on tomato (*Solanum lycopersicum*) fruit yield and quality with special emphasis on sugars, acids, ascorbate, carotenoids, and phenolic compounds. *Journal of Agricultural and Food Chemistry*, 57, 4112–4123.
- Bernstein, P.S., Li, B., Vachali, P.P., Gorusupudi, A., Shyam, R., Henriksen, B.S. et al. (2016) Lutein, zeaxanthin, and meso-zeaxanthin: the basic and clinical science underlying carotenoid-based nutritional interventions against ocular disease. *Progress in Retinal and Eye Research*, 50, 34–66.
- Borghesi, E., González-Miret, M.L., Escudero-Gilete, M.L., Malorgio, F., Heredia, F.J. & Meléndez-Martínez, A.J. (2011) Effects of salinity stress on carotenoids, anthocyanins, and color of diverse tomato genotypes. *Journal of Agricultural and Food Chemistry*, 59, 11676–11682.
- Brandt, S., Pék, Z., Barna, É., Lugasi, A. & Helyes, L. (2006) Lycopene content and colour of ripening tomatoes as affected by environmental conditions. *Journal of the Science of Food and Agriculture*, 86, 568–572.
- Bucchetti, B., Matthews, M.A., Falginella, L., Peterlunger, E. & Castellarin, S.D. (2011) Effect of water deficit on Merlot grape tannins and anthocyanins across four seasons. *Scientia Horticulturae*, 128, 297–305.
- Butelli, E., Licciardello, C., Zhang, Y., Liu, J., Mackay, S., Bailey, P. et al. (2012) Retrotransposons control fruit-specific, cold-dependent accumulation of anthocyanins in blood oranges. *The Plant Cell*, 24, 1242–1255.
- Carpentier, S., Knaus, M. & Suh, M. (2009) Associations between lutein, zeaxanthin, and age-related macular degeneration: an overview. *Critical Reviews in Food Science and Nutrition*, 49, 313–326.
- Castellarin, S.D., Matthews, M.A., Di Gasparo, G. & Gambetta, G.A. (2007) Water deficits accelerate ripening and induce changes in gene expression regulating flavonoid biosynthesis in grape berries. *Planta*, 227, 101–112.
- Cazzonelli, C.I. & Pogson, B.J. (2010) Source to sink: regulation of carotenoid biosynthesis in plants. *Trends in Plant Science*, 15, 266–274.
- Challinor, A.J., Watson, J., Lobell, D.B., Howden, S.M., Smith, D.R. & Chhetri, N. (2014) A meta-analysis of crop yield under climate change and adaptation. *Nature Climate Change*, 4, 287–291.
- Chen, Z., Yu, L., Liu, W., Zhang, J., Wang, N. & Chen, X. (2021) Research progress of fruit color development in apple (*Malus domestica* Borkh.). *Plant Physiology and Biochemistry*, 162, 267–279.
- Cheok, A., Xu, Y., Zhang, Z., Caton, P.W. & Rodriguez-Mateos, A. (2022) Betalain-rich dragonfruit (pitaya) consumption improves vascular function in men and women: adouble-blind, randomized controlled crossover trial. *The American Journal of Clinical Nutrition*, 115, 1418–1431.
- Connor, A.M., Luby, J.J., Tong, C.B.S., Finn, C.E. & Hancock, J.F. (2002) Genotypic and environmental variation in antioxidant activity, total phenolic content, and anthocyanin content among blueberry cultivars. *Journal of the American Society for Horticultural Science*, 127, 89–97.
- Cunningham, F.X. & Gantt, E. (1998) Genes and enzymes of carotenoid biosynthesis in plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, 49, 557–583.
- D'Amelia, V., Aversano, R., Chiaiese, P. & Carputo, D. (2018) The antioxidant properties of plant flavonoids: their exploitation by molecular plant breeding. *Phytochemistry Reviews*, 17, 611–625.
- Das, K. & Roychoudhury, A. (2014) Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Frontiers in environmental science*, 2, 53.
- Davies, K.M., Albert, N.W., Zhou, Y. & Schwinn, K.E. (2018) Functions of flavonoid and betalain pigments in abiotic stress tolerance in plants. *Annual Plant Reviews online*, 1, 21–62.
- Davies, K.M., Landi, M., van Klink, J.W., Schwinn, K.E., Brummell, D.A., Albert, N.W. et al. (2022) Evolution and function of redpigmentation in land plants. *Annals of Botany*, 130, 613–636.
- Davuluri, G.R., van Tuinen, A., Fraser, P.D., Manfredonia, A., Newman, R., Burgess, D. et al. (2005) Fruit-specific RNAi-mediated suppression of DET1 enhances carotenoid and flavonoid content in tomatoes. *Nature Biotechnology*, 23, 890–895.
- De, P., Martino, A., Raimondi, G. & Maggio, A. (2007) Comparative analysis of water and salt stress-induced modifications of quality parameters in cherry tomatoes. *The Journal of Horticultural Science and Biotechnology*, 82, 283–289.
- Decros, G., Baldet, P., Beauvoit, B., Stevens, R., Flandin, A., Colombié, S. et al. (2019) Get the balance right: ROS homeostasis and redox signalling in fruit. *Frontiers in Plant Science*, 10, 1091.
- Deluc, L.G., Quilici, D.R., Decendit, A., Grimplet, J., Wheatley, M.D., Schlauch, K.A. et al. (2009) Water deficit alters differentially metabolic pathways affecting important flavor and quality traits in grape berries of cabernet sauvignon and chardonnay. *BMC Genomics*, 10, 212.
- Dias, M.G., Borge, G.I.A., Kijak, K., Mandić, A.I., Mapelli-Brahm, P., Olmedilla-Alonso, B. et al. (2021) European database of carotenoid levels in foods. factors affecting carotenoid content. *Foods*, 10, 912.

- Dias, M.G., Camões, M.F.G.F.C. & Oliveira, L. (2009) Carotenoids in traditional Portuguese fruits and vegetables. *Food Chemistry*, 113, 808–815.
- Downey, M.O., Dokoozlian, N.K. & Krstic, M.P. (2006) Cultural practice and environmental impacts on the flavonoid composition of grapes and wine: a review of recent research. *American Journal of Enology and Viticulture*, 57, 257–268.
- Espley, R.V., Butts, C.A., Laing, W.A., Martell, S., Smith, H., McGhie, T.K. et al. (2014) Dietary flavonoids from modified apple reduce inflammation markers and modulate gut microbiota in mice. *The Journal of Nutrition*, 144, 146–154.
- Fanciullino, A.-L., Bidel, L.P. & Urban, L. (2014) Carotenoid responses to environmental stimuli: integrating redox and carbon controls into a fruit model. *Plant, Cell & Environment*, 37, 273–289.
- Fang, H., Dong, Y., Yue, X., Hu, J., Jiang, S., Xu, H. et al. (2019) The B-box zinc finger protein MdBBX20 integrates anthocyanin accumulation in response to ultraviolet radiation and low temperature. *Plant, Cell & Environment*, 42, 2090–2104.
- Favati, F., Lovelli, S., Galgano, F., Miccolis, V., Di Tommaso, T. & Candido, V. (2009) Processing tomato quality as affected by irrigation scheduling. *Scientia Horticulturae*, 122, 562–571.
- Fenn, M.A. & Giovannoni, J.J. (2021) Phytohormones in fruit development and maturation. *The Plant Journal*, 105, 446–458.
- Foyer, C.H. (2018) Reactive oxygen species, oxidative signaling and the regulation of photosynthesis. *Environmental and Experimental Botany*, 154, 134–142.
- Frank, H.A. & Cogdell, R.J. (1996) Carotenoids in photosynthesis. *Photochemistry and Photobiology*, 63, 257–264.
- Fredes, C., Yousef, G.G., Robert, P., Grace, M.H., Lila, M.A., Gómez, M. et al. (2014) Anthocyanin profiling of wild maqui berries (*Aristotelia chilensis*[Mol.] Stuntz) from different geographical regions in Chile: anthocyanin profiling of wild maqui berries. *Journal of the Science of Food and Agriculture*, 94, 2639–2648.
- Fu, C., Han, Y., Fan, Z., Chen, J., Chen, W., Lu, W. et al. (2016) The papaya transcription factor CpNAC1 modulates carotenoid biosynthesis through activating phytoene desaturase genes CpPDS2/4 during fruit ripening. *Journal of Agricultural and Food Chemistry*, 64, 5454–5463.
- Fu, C., Han, Y., Kuang, J., Chen, J. & Lu, W. (2017) Papaya CpEIN3a and CpNAC2 co-operatively regulate carotenoid biosynthesis-related genes CpPDS2/4, CpLCY-e and CpCHY-b during fruit ripening. *Plant and Cell Physiology*, 58, 2155–2165.
- Gambetta, G.A., Herrera, J.C., Dayer, S., Feng, Q., Hochberg, U. & Castellarin, S.D. (2020) The physiology of drought stress in grapevine: towards an integrative definition of drought tolerance. *Journal of Experimental Botany*, 71, 4658–4676.
- Gashu, K., Song, C., Dubey, A.K., Acuña, T., Sagí, M., Agam, N. et al. (2022) The effect of topo-climate variation on the secondary metabolism of berries in White grapevine varieties (*Vitis vinifera*). *Frontiers in Plant Science*, 13, 487.
- Giuliano, G. (2014) Plant carotenoids: genomics meets multi-gene engineering. *Current Opinion in Plant Biology*, 19, 111–117.
- Giuliano, G. (2017) Provitamin A biofortification of crop plants: a gold rush with many miners. *Current Opinion in Biotechnology*, 44, 169–180.
- Giuntini, D., Graziani, G., Lercari, B., Fogliano, V., Soldatini, G.F. & Ranieri, A. (2005) Changes in carotenoid and ascorbic acid contents in fruits of different tomato genotypes related to the depletion of UV-B radiation. *Journal of Agricultural and Food Chemistry*, 53, 3174–3181.
- Gonçalves, A.C., Nunes, A.R., Falcão, A., Alves, G. & Silva, L.R. (2021) Dietary effects of anthocyanins in human health: a comprehensive review. *Pharmaceuticals*, 14, 690.
- Gong, J., Zeng, Y., Meng, Q., Guan, Y., Li, C., Yang, H. et al. (2021) Red light-induced kumquat fruit coloration is attributable to increased carotenoid metabolism regulated by FcrNAC22. *Journal of Experimental Botany*, 72, 6274–6290.
- González-de-Peredo, A.V., Vázquez-Espinosa, M., Espada-Bellido, E., Ferreira-González, M., Amores-Arocha, A., Palma, M. et al. (2019) Discrimination of myrtle ecotypes from different geographic areas according to their morphological characteristics and anthocyanins composition. *Plants*, 8, 328.
- Groh, I.A.M., Riva, A., Braun, D., Sutherland, H.G., Williams, O., Bakuradze, T. et al. (2020) Long-term consumption of anthocyanin-rich fruit juice: impact on gut microbiota and antioxidant markers in lymphocytes of healthy males. *Antioxidants*, 10, 27.
- He, J., Yao, L., Pecoraro, L., Liu, C., Wang, J., Huang, L. et al. (2022) Cold stress regulates accumulation of flavonoids and terpenoids in plants by phytohormone, transcription process, functional enzyme, and epigenetics. *Critical Reviews in Biotechnology*, 43, 1–18.
- Henry-Kirk, R.A., Plunkett, B., Hall, M., McGhie, T., Allan, A.C., Wargent, J.J. et al. (2018) Solar UV light regulates flavonoid metabolism in apple (*Malus x domestica*). *Plant, Cell & Environment*, 41, 675–688.
- Hernández, V., Hellín, P., Fenoll, J. & Flores, P. (2020) Impact of nitrogen supply limitation on tomato fruit composition. *Scientia Horticulturae*, 264, 109173.
- Hochberg, U., Degu, A., Cramer, G.R., Rachmilevitch, S. & Fait, A. (2015) Cultivar specific metabolic changes in grapevines berry skins in relation to deficit irrigation and hydraulic behavior. *Plant Physiology and Biochemistry*, 88, 42–52.
- Husain, T., Fatima, A., Suhel, M., Singh, S., Sharma, A., Prasad, S.M. et al. (2020) A brief appraisal of ethylene signaling under abiotic stress in plants. *Plant Signaling & Behavior*, 15, 1782051.
- Ilić, Z.S. & Fallik, E. (2017) Light quality manipulation improves vegetable quality at harvest and postharvest: a review. *Environmental and Experimental Botany*, 139, 79–90.
- Jaakola, L. (2013) New insights into the regulation of anthocyanin biosynthesis in fruits. *Trends in Plant Science*, 18(9), 477–483.
- Jaakola, L. & Hohtola, A. (2010) Effect of latitude on flavonoid biosynthesis in plants. *Plant, Cell & Environment*, 33, 1239–1247.
- Jaakola, L., Poole, M., Jones, M.O., Kämäräinen-Karppinen, T., Koskimäki, J.J., Hohtola, A. et al. (2010) A SQUAMOSA MADS box gene involved in the regulation of anthocyanin accumulation in bilberry fruits. *Plant Physiology*, 153, 1619–1629.
- Jaleel, C.A., Manivannan, P., Wahid, A., Farooq, M., Al-Juburi, H.J., Somasundaram, R. et al. (2009) Drought stress in plants: a review on morphological characteristics and pigments composition. *International Journal of Agriculture and Biology*, 11, 100–105.
- Jezek, M., Zörb, C., Merkt, N. & Geilfus, C.-M. (2018) Anthocyanin management in fruits by fertilization. *Journal of Agricultural and Food Chemistry*, 66, 753–764.
- Jia, H.-F., Chai, Y.-M., Li, C.-L., Lu, D., Luo, J.-J., Qin, L. et al. (2011) Abscisic acid plays an important role in the regulation of strawberry fruit ripening. *Plant Physiology*, 157, 188–199.
- Jiang, W., Li, N., Zhang, D., Meinhardt, L., Cao, B., Li, Y. et al. (2020) Elevated temperature and drought stress significantly affect fruit quality and activity of anthocyanin-related enzymes in jujube (*Ziziphus jujuba* Mill. cv. 'Lingwuchangzao'). *PLoS One*, 15, e0241491.
- Joubert, C., Young, P.R., Eyéghé-Bickong, H.A. & Vivier, M.A. (2016) Field-grown grapevine berries use carotenoids and the associated xanthophyll cycles to acclimate to UV exposure differentially in high and low light (shade) conditions. *Frontiers in Plant Science*, 7, 786.
- Jucá, M.M., Cysne Filho, F.M.S., de Almeida, J.C., Mesquita, D.S., Barriga, J.R.M., Dias, K.C.F. et al. (2020) Flavonoids: biological activities and therapeutic potential. *Natural Product Research*, 34, 692–705.
- Kai, H. & Iba, K. (2014) Temperature stress in plants. In: *Encyclopedia of Life Sciences*. John Wiley & Sons, Ltd. <https://doi.org/10.1002/9780470015902.a0001320.pub2>

- Kalt, W., Cassidy, A., Howard, L.R., Krikorian, R., Stull, A.J., Tremblay, F. et al. (2020) Recent research on the health benefits of blueberries and their anthocyanins. *Advances in Nutrition*, 11, 224–236.
- Karppinen, K., Lafferty, D.J., Albert, N.W., Mikkola, N., McGhie, T., Allan, A.C. et al. (2021) MYBA and MYBPA transcription factors co-regulate anthocyanin biosynthesis in blue-coloured berries. *New Phytologist*, 232, 1350–1367.
- Karppinen, K., Tegelberg, P., Häggman, H. & Jaakola, L. (2018) Abscisic acid regulates anthocyanin biosynthesis and gene expression associated with cell wall modification in ripening bilberry (*Vaccinium myrtillus* L.) fruits. *Frontiers in Plant Science*, 9, 1259.
- Kay, C.D., Pereira-Caro, G., Ludwig, I.A., Clifford, M.N. & Crozier, A. (2017) Anthocyanins and flavanones are more bioavailable than previously perceived: a review of recent evidence. *Annual Review of Food Science and Technology*, 8, 155–180.
- Kimura, M., Rodriguez-Amaya, D. & Yokoyama, S. (1991) Cultivar differences and geographic effects on the carotenoid composition and vitamin A value of papaya. *Lebensmittel-Wissenschaft + Technologie*, 24, 415–418.
- Kliwer, W.M. (1977) Influence of temperature, solar radiation and nitrogen on coloration and composition of emperor grapes. *American Journal of Enology and Viticulture*, 28, 96–103.
- Klunklin, W. & Savage, G. (2017) Effect on quality characteristics of tomatoes grown under well-watered and drought stress conditions. *Foods*, 6, 56.
- Kokalj, D., Zlatić, E., Cigić, B., Kobav, M.B. & Vidrih, R. (2019a) Postharvest flavonol and anthocyanin accumulation in three apple cultivars in response to blue-light-emitting diode light. *Scientia Horticulturae*, 257, 108711.
- Kokalj, D., Zlatić, E., Cigić, B. & Vidrih, R. (2019b) Postharvest light-emitting diode irradiation of sweet cherries (*Prunus avium* L.) promotes accumulation of anthocyanins. *Postharvest Biology and Technology*, 148, 192–199.
- Korekar, G., Dolkar, P., Singh, H., Srivastava, R.B. & Stobdan, T. (2014) Variability and the genotypic effect on antioxidant activity, total phenolics, carotenoids and ascorbic acid content in seventeen natural population of seabuckthorn (*Hippophae rhamnoides* L.) from trans-himalaya. *LWT-Food Science and Technology*, 55, 157–162.
- Koyama, K., Ikeda, H., Poudel, P.R. & Goto-Yamamoto, N. (2012) Light quality affects flavonoid biosynthesis in young berries of cabernet sauvignon grape. *Phytochemistry*, 78, 54–64.
- Kumar, R., Khurana, A. & Sharma, A.K. (2013) Role of plant hormones and their interplay in development and ripening of fleshy fruits. *Journal of Experimental Botany*, 65, 4561–4575.
- Lado, J., Rodrigo, M.J., Cronje, P. & Zacarias, L. (2015) Involvement of lycopene in the induction of tolerance to chilling injury in grapefruit. *Postharvest Biology and Technology*, 100, 176–186.
- Lafferty, D.J., Espley, R.V., Deng, C.H., Günther, C.S., Plunkett, B. & Turner, J.L. et al. (2021) Hierarchical regulation of MYBPA1 by anthocyanin-and proanthocyanidin-related MYB proteins is conserved in vaccinium species. *Journal of Experimental Botany*, 73, 1344–1356.
- LaFountain, A.M. & Yuan, Y.W. (2021) Repressors of anthocyanin biosynthesis. *New Phytologist*, 231, 933–949.
- Lätti, A.K., Jaakola, L., Riihinen, K.R. & Kainulainen, P.S. (2010) Anthocyanin and flavonol variation in bog bilberries (*Vaccinium uliginosum* L.) in Finland. *Journal of Agricultural and Food Chemistry*, 58, 427–433.
- Lätti, A.K., Riihinen, K.R. & Kainulainen, P.S. (2008) Analysis of anthocyanin variation in wild populations of bilberry (*Vaccinium myrtillus* L.) in Finland. *Journal of Agricultural and Food Chemistry*, 56, 190–196.
- Leermakers, E.T., Darweesh, S.K., Baena, C.P., Moreira, E.M., Melo van Lent, D., Tielemans, M.J. et al. (2016) The effects of lutein on cardiometabolic health across the life course: a systematic review and meta-analysis. *The American Journal of Clinical Nutrition*, 103, 481–494.
- Leng, P., Yuan, B. & Guo, Y. (2013) The role of abscisic acid in fruit ripening and responses to abiotic stress. *Journal of Experimental Botany*, 65, 4577–4588.
- Li, G., Meng, X., Zhu, M. & Li, Z. (2019) Research progress of betalain in response to adverse stresses and evolutionary relationship compared with anthocyanin. *Molecules*, 24, 3078.
- Li, Y., Wang, H., Zhang, Y. & Martin, C. (2018) Can the world's favorite fruit, tomato, provide an effective biosynthetic chassis for high-value metabolites? *Plant Cell Reports*, 37, 1443–1450.
- Lila, M.A., Burton-Freeman, B., Grace, M. & Kalt, W. (2016) Unraveling anthocyanin bioavailability for human health. *Annual Review of Food Science and Technology*, 7, 375–393.
- Lillo, C., Lea, U.S. & Ruoff, P. (2008) Nutrient depletion as a key factor for manipulating gene expression and product formation in different branches of the flavonoid pathway. *Plant, Cell & Environment*, 31, 587–601.
- Lindbergh, C.A., Renzi-Hammond, L.M., Hammond, B.R., Terry, D.P., Mewborn, C.M., Puente, A.N. et al. (2018) Lutein and zeaxanthin influence brain function in older adults: a randomized controlled trial. *Journal of the International Neuropsychological Society*, 24, 77–90.
- Lin-Wang, K., McGhie, T.K., Wang, M., Liu, Y., Warren, B., Storey, R. et al. (2014) Engineering the anthocyanin regulatory complex of strawberry (*Fragaria vesca*). *Frontiers in Plant Science*, 5, 1–14.
- Lin-Wang, K., Micheletti, D., Palmer, J., Volz, R., Lozano, L., Espley, R. et al. (2011) High temperature reduces apple fruit colour via modulation of the anthocyanin regulatory complex. *Plant, Cell & Environment*, 34, 1176–1190.
- Liso, M., De Santis, S., Scarano, A., Verna, G., Dicarlo, M., Galleggiante, V. et al. (2018) A bronze-tomato enriched diet affects the intestinal microbiome under homeostatic and inflammatory conditions. *Nutrients*, 10, 1862.
- Lobell, D.B. & Gourdj, S.M. (2012) The influence of climate change on global crop productivity. *Plant Physiology*, 160, 1686–1697.
- Lopez, M., Candela, M.E. & Sabater, F. (1986) Carotenoids from capsicum annum fruits: influence of spectral quality of radiation. *Biologia Plantarum*, 28, 100–104.
- Lu, S., Zhang, Y., Zhu, K., Yang, W., Ye, J., Chai, L. et al. (2018) The citrus transcription factor CsMADS6 modulates carotenoid metabolism by directly regulating carotenogenic genes. *Plant Physiology*, 176, 2657–2676.
- Ma, G., Zhang, L., Kato, M., Yamawaki, K., Kiriwa, Y., Yahata, M. et al. (2012) Effect of blue and red LED light irradiation on β -cryptoxanthin accumulation in the flavedo of citrus fruits. *Journal of Agricultural and Food Chemistry*, 60, 197–201.
- Mao, W., Han, Y., Chen, Y., Sun, M., Feng, Q., Li, L. et al. (2022) Low temperature inhibits anthocyanin accumulation in strawberry fruit by activating FvMAPK3-induced phosphorylation of FvMYB10 and degradation of chalcone synthase 1. *The Plant Cell*, 34, 1226–1249.
- Maoka, T. (2020) Carotenoids as natural functional pigments. *Journal of Natural Medicines*, 74, 1–16.
- Martin, C. & Li, J. (2017) Medicine is not health care, food is health care: plant metabolic engineering, diet and human health. *New Phytologist*, 216, 699–719.
- Martinussen, I., Uleberg, E., McDougall, G.J., Stewart, D. & Junntila, O. (2010) Development and quality of cloudberry (*Rubus chamaemorus* L.) as affected by female parent, male parent and temperature. *Journal of Berry Research*, 1, 91–101.
- Medina-Puche, L., Cumplido-Laso, G., Amil-Ruiz, F., Hoffmann, T., Ring, L., Rodríguez-Franco, A. et al. (2013) MYB10 plays a major role in the regulation of flavonoid/phenylpropanoid metabolism during ripening of fragaria \times ananassa fruits. *Journal of Experimental Botany*, 65, 401–417.
- Medina-Puche, L., Cumplido-Laso, G., Amil-Ruiz, F., Hoffmann, T., Ring, L., Rodríguez-Franco, A. et al. (2014) MYB10 plays a major role in the

- regulation of flavonoid/phenylpropanoid metabolism during ripening of fragaria× ananassa fruits. *Journal of Experimental Botany*, 65, 401–417.
- Mein, J.R., Lian, F. & Wang, X.-D. (2008) Biological activity of lycopene metabolites: implications for cancer prevention. *Nutrition Reviews*, 66, 667–683.
- Mercadante, A.Z. & Rodriguez-Amaya, D.B. (1998) Effects of ripening, cultivar differences, and processing on the carotenoid composition of mango. *Journal of Agricultural and Food Chemistry*, 46, 128–130.
- Merzlyak, M.N. & Chivkunova, O.B. (2000) Light-stress-induced pigment changes and evidence for anthocyanin photoprotection in apples. *Journal of Photochemistry and Photobiology, B: Biology*, 55, 155–163.
- Mesa, T., Polo, J., Arabia, A., Caselles, V. & Munné-Bosch, S. (2022) Differential physiological response to heat and cold stress of tomato plants and its implication on fruit quality. *Journal of Plant Physiology*, 268, 153581.
- Miao, L., Zhang, Y., Yang, X., Xiao, J., Zhang, H., Zhang, Z. et al. (2016) Colored light-quality selective plastic films affect anthocyanin content, enzyme activities, and the expression of flavonoid genes in strawberry (*Fragaria×ananassa*) fruit. *Food Chemistry*, 207, 93–100.
- Mittler, R. (2002) Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science*, 7, 405–410.
- Moise, A.R., Al-Babili, S. & Wurtzel, E.T. (2014) Mechanistic aspects of carotenoid biosynthesis. *Chemical Reviews*, 114, 164–193.
- Movahed, N., Pastore, C., Cellini, A., Allegro, G., Valentini, G., Zenoni, S. et al. (2016) The grapevine VviPrx31 peroxidase as a candidate gene involved in anthocyanin degradation in ripening berries under high temperature. *Journal of Plant Research*, 129, 513–526.
- Müller, M. (2021) Foes or friends: ABA and ethylene interaction under abiotic stress. *Plants*, 10, 448.
- Ntemiri, A., Ghosh, T.S., Gheller, M.E., Tran, T.T.T., Blum, J.E., Pellanda, P. et al. (2020) Whole blueberry and isolated polyphenol-rich fractions modulate specific gut microbes in an in vitro colon model and in a pilot study in human consumers. *Nutrients*, 12, 2800.
- Osakabe, Y., Osakabe, K., Shinozaki, K. & Tran, L.S.P. (2014) Response of plants to water stress. *Frontiers in Plant Science*, 5, 5.
- Ou, J.L.S., Yang, D. & Liu, M.H. (2020) Effects of anthocyanins in composite meals on cardiometabolic Outcomes—A systematic review of randomized controlled feeding trials. *Nutrients*, 12, 3781.
- Pan, Z., Liu, Q., Yun, Z., Guan, R., Zeng, W., Xu, Q. et al. (2009) Comparative proteomics of a lycopene-accumulating mutant reveals the important role of oxidative stress on carotenogenesis in sweet Orange (*Citrus sinensis* [L.] osbeck). *Proteomics*, 9, 5455–5470.
- de Pascual-Teresa, S. & Sanchez-Ballesta, M.T. (2008) Anthocyanins: from plant to health. *Phytochemistry Reviews*, 7, 281–299.
- Patanè, C., Siah, S., Pellegrino, A., Cosentino, S.L. & Siracusa, L. (2021) Fruit yield, polyphenols, and carotenoids in long Shelf-Life tomatoes in response to drought stress and rewatering. *Agronomy*, 11, 1943.
- Perin, E.C., da Silva Messias, R., Borowski, J.M., Crizel, R.L., Schott, I.B., Carvalho, I.R. et al. (2019) ABA-dependent salt and drought stress improve strawberry fruit quality. *Food Chemistry*, 271, 516–526.
- Plunkett, B.J., Espley, R.V., Dare, A.P., Warren, B.A.W., Grierson, E.R.P., Cordiner, S. et al. (2018) MYBA from blueberry (*Vaccinium* Section *Cyanococcus*) is a subgroup 6 type R2R3MYB transcription factor that activates anthocyanin production. *Frontiers in Plant Science*, 9, 1–14.
- Poiroux-Gonord, F., Bidet, L.P.R., Fanciullino, A.-L., Gautier, H., Lauri-Lopez, F. & Urban, L. (2010) Health benefits of vitamins and secondary metabolites of fruits and vegetables and prospects to increase their concentrations by agronomic approaches. *Journal of Agricultural and Food Chemistry*, 58, 12065–12082.
- Poiroux-Gonord, F., Santini, J., Fanciullino, A.L., Lopez-Lauri, F., Giannettini, J., Sallanon, H. et al. (2013) Metabolism in Orange fruits is driven by photooxidative stress in the leaves. *Physiologia Plantarum*, 149, 175–187.
- Pojer, E., Mattivi, F., Johnson, D. & Stockley, C.S. (2013) The case for anthocyanin consumption to promote human health: a review. *Comprehensive Reviews in Food Science and Food Safety*, 12, 483–508.
- Pola, W., Sugaya, S. & Photchanachai, S. (2019) Color development and phytochemical changes in mature Green chili (*Capsicum annum* L.) exposed to red and blue light-emitting diodes. *Journal of Agricultural and Food Chemistry*, 68, 59–66.
- Qu, Z. & Zhou, G. (2016) Possible impact of climate change on the quality of apples from the major producing areas of China. *Atmosphere*, 7, 113.
- Rahim, M.A., Busatto, N. & Trainotti, L. (2014) Regulation of anthocyanin biosynthesis in peach fruits. *Planta*, 240, 913–929.
- Ravaglia, D., Espley, R.V., Henry-Kirk, R.A., Andreotti, C., Ziosi, V., Hellens, R.P. et al. (2013) Transcriptional regulation of flavonoid biosynthesis in nectarine (*Prunus persica*) by a set of R2R3 MYB transcription factors. *BMC Plant Biology*, 13, 68.
- Rehman, R.N.U., You, Y., Zhang, L., Goudia, B.D., Khan, A.R., Li, P. et al. (2017) High temperature induced anthocyanin inhibition and active degradation in malus profusion. *Frontiers in Plant Science*, 8, 1401.
- Ripoll, J., Urban, L., Staudt, M., Lopez-Lauri, F., Bidet, L.P.R. & Bertin, N. (2014) Water shortage and quality of fleshy fruits—making the most of the unavoidable. *Journal of Experimental Botany*, 65, 4097–4117.
- Rodriguez-Amaya, D.B. (2001) *A guide to carotenoid analysis in foods*, vol. 71, Washington: ILSI press.
- Rodriguez-Concepcion, M., Avalos, J., Bonet, M.L., Boronat, A., Gomez-Gomez, L., Hornero-Mendez, D. et al. (2018) A global perspective on carotenoids: metabolism, biotechnology, and benefits for nutrition and health. *Progress in Lipid Research*, 70, 62–93.
- Rohloff, J., Uleberg, E., Nes, A., Krogstad, T., Nestby, R. & Martinussen, I. (2015) Nutritional composition of bilberries (*Vaccinium myrtillus* L.) from forest fields in Norway—Effects of geographic origin, climate, fertilization and soil properties. *Journal of Applied Botany and Food Quality*, 88, 274–287.
- Rosalie, R., Léchaudel, M., Dhuique-Mayer, C., Dufossé, L. & Joas, J. (2018) Antioxidant and enzymatic responses to oxidative stress induced by cold temperature storage and ripening in mango (*Mangifera indica* L. cv. 'Cogshall') in relation to carotenoid content. *Journal of Plant Physiology*, 224–225, 75–85.
- Ryu, S., Han, J.H., Cho, J.G., Jeong, J.H., Lee, S.K. & Lee, H.J. (2020) High temperature at veraison inhibits anthocyanin biosynthesis in berry skins during ripening in 'kyoho' grapevines. *Plant Physiology and Biochemistry*, 157, 219–228.
- Saini, R.K. & Keum, Y.-S. (2018) Significance of genetic, environmental, and pre-and postharvest factors affecting carotenoid contents in crops: a review. *Journal of Agricultural and Food Chemistry*, 66, 5310–5324.
- Samkumar, A., Jones, D., Karppinen, K., Dare, A.P., Sipari, N., Espley, R.V. et al. (2021) Red and blue light treatments of ripening bilberry fruits reveal differences in signalling through abscisic acid-regulated anthocyanin biosynthesis. *Plant, Cell & Environment*, 44, 3227–3245.
- Samkumar, A., Karppinen, K., McGhie, T.K., Espley, R.V., Martinussen, I. & Jaakola, L. (2022) Flavonoid biosynthesis is differentially altered in detached and attached ripening bilberries in response to spectral light quality. *Frontiers in Plant Science*, 13, 2529.
- Santesteban, L.G., Miranda, C. & Royo, J.B. (2011) Regulated deficit irrigation effects on growth, yield, grape quality and individual anthocyanin composition in *Vitis vinifera* L. cv. 'tempranillo'. *Agricultural Water Management*, 98, 1171–1179.
- Schertl, P. & Braun, H.-P. (2014) Respiratory electron transfer pathways in plant mitochondria. *Frontiers in Plant Science*, 5, 163.
- Sicilia, A., Scialò, E., Puglisi, I. & Lo Piero, A.R. (2020) Anthocyanin biosynthesis and DNA methylation dynamics in sweet Orange fruit

- [*Citrus sinensis* L. (Osbeck)] under cold stress. *Journal of Agricultural and Food Chemistry*, 68, 7024–7031.
- Sikuten, I., Stambuk, P., Tomaz, I., Marchal, C., Kontic, J.K., Lacombe, T. et al. (2021) Discrimination of genetic and geographical groups of grape varieties (*Vitis vinifera* L.) based on their polyphenolic profiles. *Journal of Food Composition and Analysis*, 102, 104062.
- Sivankalyani, V., Feygenberg, O., Diskin, S., Wright, B. & Alkan, N. (2016) Increased anthocyanin and flavonoids in mango fruit peel are associated with cold and pathogen resistance. *Postharvest Biology and Technology*, 111, 132–139.
- Stanley, L. & Yuan, Y.-W. (2019) Transcriptional regulation of carotenoid biosynthesis in plants: so many regulators, so little consensus. *Frontiers in Plant Science*, 10, 1017.
- Stracke, R., Werber, M. & Weisshaar, B. (2001) The R2R3-MYB gene family in *Arabidopsis thaliana*. *Current Opinion in Plant Biology*, 4, 447–456.
- Sudheeran, P., Feygenberg, O., Maurer, D. & Alkan, N. (2018) Improved cold tolerance of mango fruit with enhanced anthocyanin and flavonoid contents. *Molecules*, 23, 1832.
- Sun, X., Jia, X., Huo, L., Che, R., Gong, X., Wang, P. et al. (2018) MdATG18a overexpression improves tolerance to nitrogen deficiency and regulates anthocyanin accumulation through increased autophagy in transgenic apple. *Plant, Cell & Environment*, 41, 469–480.
- Sun, X., Li, X., Wang, Y., Xu, J., Jiang, S. & Zhang, Y. (2022) MdMKK9-mediated the regulation of anthocyanin synthesis in red-fleshed apple in response to different nitrogen signals. *International Journal of Molecular Sciences*, 23, 7755.
- Tao, R., Bai, S., Ni, J., Yang, Q., Zhao, Y. & Teng, Y. (2018) The blue light signal transduction pathway is involved in anthocyanin accumulation in 'red zaosu' pear. *Planta*, 248, 37–48.
- Tapia, G., Castro, M., Gaete-Eastman, C. & Figueroa, C.R. (2022) Regulation of anthocyanin biosynthesis by drought and UV-B radiation in wild tomato (*Solanum peruvianum*) fruit. *Antioxidants*, 11, 1639.
- Thomson, G.E., Turpin, S. & Goodwin, I. (2018) A review of preharvest anthocyanin development in full red and blush cultivars of european pear. *New Zealand Journal of Crop and Horticultural Science*, 46, 81–100.
- Tian, J., Han, Z., Zhang, L., Song, T.-T., Zhang, J., Li, J.-Y. et al. (2015) Induction of anthocyanin accumulation in crabapple (*Malus cv.*) leaves by low temperatures. *HortScience*, 50, 640–649.
- Tomes, M.L. (1963) Temperature inhibition of carotene synthesis in tomato. *Botanical Gazette*, 124, 180–185.
- Tramblay, Y., Koutroulis, A., Samaniego, L., Vicente-Serrano, S.M., Volaire, F., Boone, A. et al. (2020) Challenges for drought assessment in the Mediterranean region under future climate scenarios. *Earth-Science Reviews*, 210, 103348.
- Ubi, B.E., Honda, C., Bessho, H., Kondo, S., Wada, M., Kobayashi, S. et al. (2006) Expression analysis of anthocyanin biosynthetic genes in apple skin: effect of UV-B and temperature. *Plant Science*, 170, 571–578.
- Uleberg, E., Rohloff, J., Jaakola, L., Tröst, K., Junttila, O., Häggman, H. et al. (2012) Effects of temperature and photoperiod on yield and chemical composition of Northern and Southern clones of bilberry (*Vaccinium myrtillus* L.). *Journal of Agricultural and Food Chemistry*, 60, 10406–10414.
- Ullah, A., Manghwar, H., Shaban, M., Khan, A.H., Akbar, A., Ali, U. et al. (2018) Phytohormones enhanced drought tolerance in plants: a coping strategy. *Environmental Science and Pollution Research*, 25, 33103–33118.
- Vighi, I.L., Crizel, R.L., Perin, E.C., Rombaldi, C.V. & Galli, V. (2019) Crosstalk during fruit ripening and stress response among abscisic acid, calcium-dependent protein kinase and phenylpropanoid. *Critical Reviews in Plant Sciences*, 38, 99–116.
- Wang, F., Sha, J., Chen, Q., Xu, X., Zhu, Z., Ge, S. et al. (2020) Exogenous abscisic acid regulates distribution of 13C and 15N and anthocyanin synthesis in 'RedFuji' apple fruit under high nitrogen supply. *Frontiers in Plant Science*, 10, 1738.
- Wang, T., Zhang, H. & Zhu, H. (2019) CRISPR technology is revolutionizing the improvement of tomato and other fruit crops. *Horticulture Research*, 6, 77.
- Wang, Y.-Y., Cheng, Y.-H., Chen, K.-E., & Tsay, Y.-F. (2018) Nitrate transport, signaling, and use efficiency. *Annual Review of Plant Biology*, 69, 85–122.
- West, K.P. & Darnton-Hill, I. (2008) Vitamin A deficiency. In *Nutrition and health in developing countries*. Springer. pp. 377–433.
- Wheeler, S., Loveys, B., Ford, C. & Davies, C. (2009) The relationship between the expression of abscisic acid biosynthesis genes, accumulation of abscisic acid and the promotion of *Vitis vinifera* L. berry ripening by abscisic acid. *Australian Journal of Grape and Wine Research*, 15, 195–204.
- Woznicki, T., Heide, O., Sønsteby, A., Wold, A.-B. & Remberg, S. (2015) Yield and fruit quality of black currant (*Ribes nigrum* L.) are favoured by precipitation and cool summer conditions. *Acta Agriculturae Scandinavica, Section B—Soil & Plant Science*, 65, 702–712.
- Xu, H., Wang, N., Liu, J., Qu, C., Wang, Y., Jiang, S. et al. (2017) The molecular mechanism underlying anthocyanin metabolism in apple using the MdMYB16 and MdbHLH33 genes. *Plant Molecular Biology*, 94, 149–165.
- Yamaguchi-Shinozaki, K. & Shinozaki, K. (2005) Organization of cis-acting regulatory elements in osmotic- and cold-stress-responsive promoters. *Trends in Plant Science*, 10, 88–94.
- Yamane, T., Jeong, S.T., Goto-Yamamoto, N., Koshita, Y. & Kobayashi, S. (2006) Effects of temperature on anthocyanin biosynthesis in grape berry skins. *American Journal of Enology and Viticulture*, 57, 54–59.
- Yang, B., Tang, J., Yu, Z., Khare, T., Srivastav, A., Dahir, S. et al. (2019) Light stress responses and prospects for engineering light stress tolerance in crop plants. *Journal of Plant Growth Regulation*, 38, 1489–1506.
- Yao, G., Ming, M., Allan, A.C., Gu, C., Li, L., Wu, X. et al. (2017) Map-based cloning of the pear gene MYB114 identifies an interaction with other transcription factors to coordinately regulate fruit anthocyanin biosynthesis. *The Plant Journal*, 92, 437–451.
- Yoshikawa, H., Honda, C. & Kondo, S. (2007) Effect of low-temperature stress on abscisic acid, jasmonates, and polyamines in apples. *Plant Growth Regulation*, 52, 199–206.
- Zhang, R.-X., Zhu, W.-C., Cheng, G.-X., Yu, Y.-N., Li, Q.-H., ul Haq, S. et al. (2020) A novel gene, CaATHB-12, negatively regulates fruit carotenoid content under cold stress in *Capsicum annuum*. *Food & Nutrition Research*, 64, 3729.
- Zhang, Y., Butelli, E. & Martin, C. (2014) Engineering anthocyanin biosynthesis in plants. *Current Opinion in Plant Biology*, 19, 81–90.
- Zhao, Y., Dong, W., Wang, K., Zhang, B., Allan, A.C., Lin-Wang, K. et al. (2017) Differential sensitivity of fruit pigmentation to ultraviolet light between two peach cultivars. *Frontiers in Plant Science*, 8, 1552.
- Zhou, H., Lin-Wang, K., Wang, H., Gu, C., Dare, A.P., Espley, R.V. et al. (2015) Molecular genetics of blood-fleshed peach reveals activation of anthocyanin biosynthesis by NAC transcription factors. *The Plant Journal*, 82, 105–121.
- Zhu, F., Luo, T., Liu, C., Wang, Y., Yang, H., Yang, W. et al. (2017) An R2R3-MYB transcription factor represses the transformation of α - and β -branch carotenoids by negatively regulating expression of CrBCH2 and CrNCED5 in flavedo of citrus reticulata. *New Phytologist*, 216, 178–192.
- Zhu, M., Fang, W., Chen, C., Wang, L. & Cao, K. (2021) Effects of shading by bagging on carotenoid accumulation in peach fruit flesh. *Journal of Plant Growth Regulation*, 40, 1912–1921.
- Zifkin, M., Jin, A., Ozga, J.A., Zaharia, L.I., Scherthner, J.P., Gesell, A. et al. (2012) Gene expression and metabolite profiling of developing

- highbush blueberry fruit indicates transcriptional regulation of flavonoid metabolism and activation of abscisic acid metabolism. *Plant Physiology*, 158, 200–224.
- Zoratti, L., Jaakola, L., Häggman, H. & Giongo, L. (2015) Anthocyanin profile in berries of wild and cultivated *vaccinium* spp. along altitudinal gradients in the Alps. *Journal of Agricultural and Food Chemistry*, 63, 8641–8650.
- Zoratti, L., Karppinen, K., Luengo Escobar, A., Häggman, H. & Jaakola, L. (2014) Light-controlled flavonoid biosynthesis in fruits. *Frontiers in Plant Science*, 5, 534.
- Zushi, K. & Matsuzoe, N. (1998) Effect of soil water deficit on vitamin C, sugar, organic acid, amino acid and carotene contents of large-fruited tomatoes. *Engei Gakkai Zasshi*, 67, 927–933.

How to cite this article: Espley, R. V. & Jaakola, L. (2023) The role of environmental stress in fruit pigmentation. *Plant, Cell & Environment*, 1–17. <https://doi.org/10.1111/pce.14684>